

COMPOSITIONS AND METHODS FOR PROTECTING CULTIVATED
PLANTS FROM HERBICIDAL INJURY

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority to U.S. Provisional Application Serial No. 60/092,993 which was filed on July 16, 1998.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

None.

BACKGROUND OF THE INVENTION

(1) Field of the Invention

The present invention relates to compositions comprising a herbicide, or combinations of herbicides, with or without a safener, and a repellent adjuvant, wherein the repellent adjuvant modifies the surface properties of the composition so that retention of the composition on foliage of the cultivated plant is reduced. In particular, the herbicide composition comprises a repellent adjuvant that is an aqueous solution of an alkyltrialkoxysilane such as methyltrimethoxysilane and a water soluble silane coupling agent such as N-(2-aminoethyl)-3-aminopropyltrimethoxysilane or an aqueous solution of an organosiliconate such as sodium methyl siliconate.

(2) Description of Related Art

Many herbicides will cause injury to certain crop plants when applied in amounts that are effective in controlling weed growth. The damage to crop plants can be particularly severe when the crop plant is in an early stage of development, which is precisely the time when control of weed growth is most important. For this reason many herbicides are unsuitable for controlling

weeds when crop plants are at particular stages of growth. Therefore, the inability to control weed growth results in lower crop yield and reduced crop quality because the weeds compete with the crop plant for nutrients, light and water. In an attempt to broaden the usefulness of various herbicides, various herbicide compositions have been developed which contain compounds known as "safeners", also referred to as "antidotes", "protectants" or "antagonists."

The concept of using safeners to enhance the tolerance of crops to herbicides was developed in the late 1940's and has led to the development of particular herbicidal compositions that have the ability to control weed growth but without adversely affecting the growth or yield of a particular crop plant. However, the identification of an antidote which safens a particular class of herbicide or mixture of herbicides is not a theoretical determination but must be determined empirically. This determination is performed by observing the complex interaction of many biological and chemical factors, including the type of herbicide, the weed species to be controlled, the crop plant to be protected from weed competition and herbicidal injury, the developmental stage of the crop plant, and the safening compound itself. Moreover, the safener and herbicide must possess physico-chemical properties which allow an environmentally acceptable and stable product to be prepared. Therefore, because the discovery of herbicide/safener compositions is so empirically based, the development of effective herbicide/safener combinations is an expensive and unpredictable undertaking.

Isoxaflutole, an isoxazole herbicide, is a new soil-applied herbicide for use in corn. Researchers have reported that isoxaflutole provides excellent preemergence control of several broadleaf and annual grass weed species at rates ranging from 53 g/ha to 158

g/ha in conventional tillage and no-tillage corn (Bhowmik and Probst, Weed Sci. Soc. Am. Abstr. 36: 13 (1996); Curvey and Kapusta, North Central Weed Sci. Soc. 51: 57-58 (1996); Geier and Stahlman, North Central Weed Sci. Soc. 52: 81 (1997); Luscombe et al., Proc. North Central Weed Sci. Soc. 59: 57-58(1994); Mosier et al., Proc. North Central Weed Sci. Soc. 50: 74 (1995); Obermeier et al., Proc. North Central Weed Sci. Soc. 50: 25 (1995); Simkins et al., Proc. North central Weed Sci. Soc. 50: 25 (1995); Veilleux et al., North central Weed Sci. Soc. 50: 75 (1995); Vrabel et al., Proc. North Central Weed Sci. Soc. 24-25 (1995); Wrucke et al., Proc. North Central Weed Sci. Soc. 52: 17(1997); and Young et al., Weed Sci. Soc. Am. Abstr. 38: 8 (1998)). The mode of action of isoxaflutole is the competitive inhibition of the 4-hydroxyphenylpyruvate dioxygenase enzyme (EC 1.13.11.27) (Pallett et al., Pestic. Biochem. Physiol. 62: 113-124 (1998); Viviani et al., Pestic. Biochem. Physiol. 62: 125-134 (1998)). Inhibition of this enzyme disrupts carotenoid biosynthesis causing a bleaching symptomology in susceptible species similar to herbicides that disrupt carotenoid biosynthesis by targeting the phytoene desaturase enzyme (Luscombe and Pallett, Pestic. Outlook, 29-32 (1996); and Pallett et al., Pestic. Sci. 50: 83-84 (1997)). While isoxaflutole has been shown to be an effective herbicide for controlling weeds, isoxaflutole also has been shown to cause injury to corn and other crop plants, especially when applied postemergent. For an example, see U.S. Patent No. 5,627,131 to Shribbs et al.

Therefore, there is a need for a method for applying a herbicide post-emergence to a wide variety of cultivated plants that does not depend on the identity of the herbicide or the use of a safener. Such a composition would not be injurious to the cultivated plant but would maintain its ability to control weeds. The present invention provides compositions and methods

for their use that satisfy this need.

SUMMARY OF THE INVENTION

5 The present invention provides compositions
comprising a herbicide, or a combination of herbicides,
with or without a safener, and a repellant adjuvant
which modifies the surface properties of the composition
so that retention of the composition on foliage of the
cultivated plant is reduced. An important element of
10 the composition is the repellant adjuvant. When the
repellant adjuvant is mixed with the herbicide or
combination of herbicides, with or without a safener, to
form the composition of the present invention and
applied to the plants with a sprayer, the emitted
15 composition spray forms spherical particles which then
bounce off the plant foliage. In this manner the
repellant adjuvant prevents retention of the composition
by the plant. Thus, herbicidal injury to the cultivated
plant is reduced because the herbicide composition is
20 directed to the soil. Therefore, the present invention
is particularly useful when it comprises herbicides that
exert their primary effects at the soil.


 The present invention provides a composition
for protecting cultivated plants comprising (a) at
25 least one herbicide and (b) a repellent adjuvant for
modifying the surface properties of the composition so
that retention of the composition on foliage of the
cultivated plant is reduced. The composition of the
present invention further comprises compositions wherein
30 the herbicide is selected from the group consisting of
acetanilides, acetamides, acetolactate synthase
inhibitors, isoxazoles, diketonitriles, triketonitriles
dinitroanilines, triazines, substituted ureas,
ethofumerates, isoxafen, oxodiazon, dithiopyr and
35 combinations thereof. Further still, the composition of
the present invention comprises a composition wherein
the composition further comprises a safener. In

particular embodiments of the present invention, the safener is selected from the group consisting of MON 4660, 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azaspiro(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinoline, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

The present invention further provides a method for reducing injury to cultivated plants, by the herbicide, by applying as a spray at least one herbicide with or without a safener in a composition with the repellent adjuvant which modifies the surface properties of the composition so that retention of the composition on foliage of the cultivated plant is reduced. The present invention provides a method for protecting cultivated plants, the composition comprising (a) a t least one herbicide and (b) a repellent adjuvant for modifying the surface properties of the composition so that retention of the composition on foliage of the cultivated plant is reduced. The present invention further provides a method for using a composition wherein the herbicide is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles,

triketonitriles dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Further still, the present invention provides a method for using the composition wherein the composition further comprises a safener. In particular embodiments of the present invention, the safener is selected from the group consisting of 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidiny]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

To make the composition of the present invention or to practice the method of the present invention, it is preferable that the repellent adjuvant be selected from the group consisting of an aqueous solution of an alkali metal organosiliconate and an aqueous solution of a water soluble siloxane solution. The organosiliconate having the formula:



wherein X denotes sodium or potassium, and R is methyl, ethyl, or propyl, and the ration of Si:X is about 1:1; and an aqueous solution of a water soluble coupling

agent and an alkyltrialkoxysilane selected from the group consisting of alkyltrialkoxysilanes with C1 to C6 alky groups on silicon and a blend of alkyltrialkoxysilanes each with a C1 to C6 alkyl groups on silicon. In a preferred embodiment, the repellent adjuvant is selected from the group consisting of an aqueous solution of sodium methyl silicate and an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane.

OBJECTS

Therefore it is an object of the present invention to provide herbicide compositions that can be applied postemergence which do not injure cultivated plants while maintaining the herbicide's ability to effectively control weeds.

These and other objects will become increasingly apparent through the following detailed description of the invention and examples.

DESCRIPTION OF THE DRAWINGS

Figure 1 is a photograph that shows the ability of the composition of the present invention comprising an aqueous solution of sodium methyl silicate to prevent corn injury. The corn plants were treated from left to right: untreated control, corn treated with isoxaflutole in combination with DUAL II (metolachlor and benoxacor), and corn treated with a composition of the present invention (a combination of isoxaflutole, DUAL II, and sodium methyl silicate). The sodium methyl silicate was applied at a rate 0.25%. The photograph shows the corn 8 days after treatment.


Figure 2 is a photograph that shows the ability of the composition of the present invention comprising an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane

to prevent corn injury. The corn plants were treated from left to right: untreated control, corn treated with a combination of isoxaflutole and DUAL II, and corn treated with a composition of the present invention (a combination of isoxaflutole, DUAL II, and an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane). The aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane was applied at a rate 0.25%. The photograph shows the corn 8 days after treatment.

DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention provides a composition for protecting cultivated plants comprising a herbicide, or combination of herbicides, with or without a safener, and a repellent adjuvant for modifying the surface properties of the composition so that retention of the composition on foliage of the cultivated plant is reduced. The herbicide comprising the present invention can be selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxadiazon, dithiopyr and combinations thereof. In particular, it is desirable that the herbicide or combination of herbicides exert their primary effects at the soil. The repellent adjuvant comprising the composition is a silicone-based aqueous solution that forms an emulsion when in combination with the herbicide, which has modifies properties. The emulsion forms spherical particles when sprayed from a sprayer. The spherical particles bounce off the foliage of the plant to the ground where the herbicide then exerts its effect. One type of repellent adjuvant suitable for use in the present invention is exemplified by an aqueous solution of the

organosiliconate having the formula:



5 wherein X denotes sodium or potassium, and R is methyl,
ethyl, or propyl, and the ratio of Si:X is about 1:1.
In a preferred embodiment of an adjuvant of this type,
the organosiliconate is sodium methyl siliconate,
potassium methyl siliconate, or a mixture thereof. In
10 a most preferred embodiment, the aqueous solution of
organosiliconate consists essentially of 32 weight
percent of sodium methyl siliconate and 67 weight
percent of water. A second type of repellent adjuvant
suitable for use in the present invention is an aqueous
15 solution of a water soluble silane coupling agent and an
alkyltrialkoxysilane, the alkyltrialkoxysilane being
selected from the group consisting of
alkyltrialkoxysilanes with C1 to C6 alkyl groups on
silicon and a blend of alkyltrialkoxysilanes each with
20 a C1 to C6 alkyl groups on silicon, the
alkyltrialkoxysilane and the silane coupling agent
preferably being present in the aqueous solution in the
mole ratio of between about 0.5:1.0 to about 3.0:1.0.
In a preferred embodiment, the water soluble silane
25 coupling agent is N-(2-aminoethyl)-3-aminopropyl-
trimethoxysilane and the alkyltrialkoxysilane is
methyltrimethoxysilane. In a most preferred embodiment,
the methyltrimethoxysilane, the N-(2-aminoethyl)-3-
aminopropyltrimethoxysilane, and the water are in an
30 aqueous solution consisting of a weight percent ratio of
35.7:58.2:6.1. Therefore, the silicon-based aqueous
solution comprising the present invention is selected
from the group consisting of an aqueous solution of an
alkali metal organosiliconate and an aqueous solution of
35 a water soluble silane coupling agent and an
alkyltrialkoxysilane.

The composition of the present invention can

further include a second herbicide, an enhancer, or an adjuvant, which increases the activity or absorption of the first herbicide. In particular, the second herbicide is an acetanilide herbicide wherein the acetanilide herbicide increases the activity or absorption of the first herbicide in the composition. An example of an acetanilide herbicide is metolachlor or a mixture of the herbicide metolachlor and the safener benoxacor. In a preferred embodiment of the invention, the composition comprises an isoxazole herbicide such as isoxaflutole, an activator herbicide such as metolachlor, and the repellent adjuvant. In a more preferred embodiment of the invention, the composition comprises an isoxazole herbicide such as isoxaflutole, an activator herbicide such as metolachlor which is in combination with a safener such as benoxacor, and the repellent adjuvant.

The activity of certain herbicides can be increased by compounds that enhance absorption of the herbicide. Therefore, the present invention further includes compositions that comprise a herbicide, the repellent adjuvant, and an enhancer adjuvant which is an oil based adjuvant. In particular, the oil-based adjuvant is selected from the group consisting of a crop oil concentrate, a free fatty acid, and an esterified and saponified oil. Examples of herbicides that are known to work well in the presence of an oil-based adjuvant are the herbicides selected from the group consisting of cyclohexanidiones, aryloxyphenoxy, imidazolinone, and sulfonylurea herbicides.

The present invention also relates to a method for protecting cultivated plants including applying a herbicide formulation that has herbicidal activity from soil, the improvement comprising using as the herbicidal formulation a homogenous aqueous dispersion of the composition comprising a herbicide, and a repellent adjuvant for modifying the surface properties the

composition so that retention of the composition on foliage of the cultivated plant is reduced. The herbicide in the method of the present invention is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. The repellent adjuvant is selected from the group consisting of an aqueous solution of an alkali metal organosiliconate and an aqueous solution of a water soluble silane coupling agent and an alkyltrialkoxysilane. In a preferred embodiment, the repellent adjuvant is selected from the group consisting of an aqueous solution of methyl siliconate and an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane. Optionally, the present invention can comprise a safener. In particular embodiments, the present invention optionally includes a safener selected from the group consisting of MON 4660, 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapiridin-4,5-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinazoline, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-

(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

5 The composition can further comprise an acetanilide herbicide as a second herbicide, which increases the activity or absorption of the first herbicide. In particular, the acetanilide herbicide in the composition can be metolachlor or a mixture of the herbicide metolachlor and the safener benoxacor. In a preferred embodiment of the invention the first herbicide comprising the composition is an isoxazole herbicide such as isoxaflutole.

10 The present invention further provides an improved method for protecting cultivated plants including the step of applying a herbicide compound, postemergence to the cultivated plants, the improvement which comprises adding a repellent adjuvant to the herbicide compound so that when the compound is applied to the cultivated plants, the compound bounces off foliage of cultivated plants to the soil. In the improved method, the repellent adjuvant is selected from any one of the repellent adjuvants disclosed herein. In the improved method, the herbicidal formulation can further comprise an enhancement material which enhances the activity of the herbicide. Enhancement materials can among other things include a second herbicide that activates or potentiates the activity of the first herbicide, or an oil-based adjuvant, or combinations of both. In a preferred embodiment, the herbicide is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Optionally, the preferred method can include a safener. In particular embodiments, the present invention optionally includes a safener selected from the group consisting of MON 4660, 2,2-dichloro-N,N-

di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-
2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-
dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-
dichloroacetyl oxazolidine, N,N-diallyl-2,2-
5 dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-
dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-
N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-
dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-
10 dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-
(dichloroacetyl)-2,2-dimethyl-5-oxalidiny]pyridine, 4-
(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-
dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-
isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-
2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-
15 tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-
diazacyclononane, 1-(dichloroacetyl)-1-
azaspiro[4,4]nonane, and combinations thereof.

The present invention further provides a
method for protecting cultivated plants without injuring
cultivated plants, the steps comprising: (a) providing
a herbicidal formulation comprising one or more
herbicides admixed with a repellent adjuvant wherein the
repellant adjuvant modifies the surface properties of
the formulation thereby reducing retention of the
20 formulation on foliage of the cultivated plants; (b) and
applying the formulation to the cultivated plants
wherein the formulation bounces off the foliage onto the
soil wherein the formulation protects the cultivated
plants without injuring the cultivated plants. In the
25 method, the repellant adjuvant is any one of the
repellant adjuvants disclosed herein. In the present
invention, the herbicidal formulation can further
comprise an enhancement material which enhances the
activity of the herbicide. In particular embodiments of
30 the present invention, the herbicide is selected from
the group consisting of acetanilides, acetamides,
acetolactate synthase inhibitors, isoxazoles,

diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Optionally, the present invention can comprise a safener. In particular embodiments, the present invention optionally comprises a safener selected from the group consisting of MON 4660, 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetylloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidiny]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinoline, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

The present invention further provides a method for inhibiting a weed without injuring turfgrass, the steps comprising (a) providing a liquid dispersion of a herbicidal formulation comprising one or more herbicides admixed with a repellent adjuvant wherein the repellent adjuvant modifies the surface properties of the formulation thereby reducing retention of the formulation on foliage of the turfgrass; and (b) applying the formulation to the crop plant wherein the formulation bounces off the foliage onto the soil wherein the formulation inhibits growth of the weed. In the method for protecting a turfgrass, the repellent adjuvant is any one of the repellent adjuvants disclosed

herein. In the method, the herbicidal formulation can further comprise an enhancement material which enhances the activity of the herbicide. In particular embodiments of the method, the herbicide is selected from the group consisting of acetanilides, acetamides, acetolactate synthase inhibitors, isoxazoles, diketonitriles, triketonitriles, dinitroanilines, triazines, substituted ureas, ethofumerates, isoxafen, oxodiazon, dithiopyr and combinations thereof. Optionally, the present invention can comprise a safener. In particular embodiments, the present invention optionally includes a safener selected from the group consisting of 2,2-dichloro-N,N-di-2-propenylacetamide, 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine, 2,2,5-trimethyl-N-dichloroacetyloxazolidine, 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, N,N-diallyl-2,2-dichloroacetamide, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapero-(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof.

The amount of herbicide comprising the composition of the present invention and used in the method of the invention varies according to a number of parameters including the cultivated plant to be protected, the weed species to be controlled, and the edaphic and climatic conditions prevailing. In general, a rate of application from about 10 to 210 grams per

hectare (g/ha) of herbicide is suitable, preferably 50 to about 158 g/ha. The rate of the repellent adjuvant in the composition can be from 0.25% to 1.0%, preferably at a rate of 0.5%.

5 According to general cultivation practices, herbicides are mixed in a tank and applied to the plants using a sprayer. The practitioner will mix various combinations of herbicides in the tank, and in some cases, will include a safener to ameliorate the herbicide's activity towards the plant to be protected. 10 In practicing the present invention, the practitioner in addition to the mixture of herbicides, with or without a safener, in the tank will include the repellent adjuvant to make the composition of the present invention. 15

Cultivated plants within the meaning of the present invention includes any plant cultivated for food or ornamentation with the exception of weeds. The cultivated plants to be protected by the method of the present invention include crop plants of which corn, 20 sugarcane, beans, rice, wheat, oats, sorghum, and a wide variety of vegetables such as tomatoes, and fruits such as strawberries are examples. In a preferred embodiment, the method of the invention is performed where the crop to be protected is corn (*Zea mays*), 25 sorghum (*Sorghum halepense*), sorghum (*Sorghum bicolor*), soybean (*Glycine max*) or dry bean (*Phaseolus vulgaris* L.). Examples of other cultivated plants that can be protected from a herbicide or combination of herbicides according to the present invention are turfgrasses; 30 flowering garden plants such as roses, tulips, carnations, orchids and the like; various herb plants such as parsley, sage, rosemary, and thyme; ornamental plants such as shrubs, holly, juniper, and spice plants.

35 Thus, the objective of the present invention is to protect cultivated plants from injury from preemergence herbicides applied postemergence to the

crop. Generally, these herbicides have activity from the soil, however it is not necessary that they have activity from the soil. The present invention results in decreased herbicide injury to the cultivated plant because it decreases foliar retention and adsorption of the herbicide by the cultivated plant. Prior to the present invention, the use of certain herbicide combinations as a postemergence application were precluded because the combination either caused injury to the cultivated plant, or caused injury to the cultivated plants because the cultivated plants were in a particularly sensitive stage at the time of the herbicide application. Therefore, many herbicide combinations cannot be used for a wide variety of cultivated plants. While the examples disclosed herein describe use of the present invention for postemergence herbicides, the present invention is not to be construed as being limited to postemergence herbicides. For example, it can be desirable to apply a preemergence pesticide to an existing turfgrass stand to kill germinating and emerging crabgrass seedlings. The present invention comprising a preemergence herbicide and a repellant adjuvant would direct the herbicide to the soil wherein the herbicide would be active.

In addition to soil activity, previous research has shown that isoxaflutole has foliar activity on a number of weed species (Sprague et al., Weed Sci. Soc. Amer. Abstr. 37: 5 (1997); Vrable et al., *ibid.*; and Young and Hart, Weed Sci. 46: 397-402 (1998)), possibly allowing for the use of postemergence applications of isoxaflutole for weed control. In fact, Sprague et al. (*ibid.*) reported that postemergence applications of isoxaflutole at 105 g/ha controlled common lambsquarters, common ragweed, redroot pigweed, and velvetleaf greater than 90% and when it was tank-mixed with metolachlor/benoxacor, the mixture controlled foxtail by greater than 80%. But this tank-mixture

greatly reduced corn tolerance when it was applied after the corn had emerged. The basis for this corn injury was the result of increased isoxaflutole retention and subsequent absorption. This result indicated that metolachlor/benoxacor may act similar to various spray adjuvants that increase herbicide retention and thereby facilitating its subsequent absorption. Spray adjuvants are normally added to foliar-applied herbicide spray solutions to maximize the effectiveness of the herbicide. These adjuvants usually exert this enhancing effect by increasing herbicide spray retention on the leaf surface and by increasing herbicide penetration into the plant cuticle. A major barrier in the retention of a herbicide is the surface tension of the spray droplets. Adjuvants such as non-ionic surfactants (NIS) and 28% urea ammonium nitrate (UAN) have been found to decrease the surface tension of spray droplets, which results in an increase in surface coverage of the spray solution (De Ruiter et al., Weed Sci. 38: 567-572 (1990); Stevens et al., Pestic. Sci. 38: 237-245 (1993)). However, there are apparently no adjuvants which can be used to modify the surface properties of a herbicide solution, which in turn decreases herbicide retention and, therefore, the herbicide absorption. The present invention provides herbicide mixtures containing adjuvants which function as repellant adjuvants because they modify the surface properties of the mixtures. It is theorized that the repellant adjuvants modify the surface property of the mixture by causing an increase in the surface tension of the mixture, which results in spray droplets of increased surface tension. Because of the increased surface tension, the herbicide spray droplets are not retained by the plant foliage. Thus, the present invention permits the postemergence application of the herbicide mixtures disclosed herein.

The herbicide isoxaflutole by itself is not

injurious to corn when applied preemergence to the corn. However, when isoxaflutole is applied to corn postemergence it causes injury to the corn. The injury is particularly severe when isoxaflutole is combined with metolachlor and benoxacor, and the combination is applied to corn plants in the spike, 2-leaf, or 4-leaf stage. The reason is that the metolachlor or any other acetanilide herbicide applied with isoxaflutole increases spray retention on the corn leaves, which ultimately increases the absorption of isoxaflutole or any other isoxazole by the corn plant. For example, DUAL II is an oily composition which when mixed with the herbicide forms a composition that facilitates absorption of the herbicide by the leaf. It is the increased absorption of the isoxazole herbicide that causes the injury to corn. Therefore, when acetanilide herbicides are used in combination with other herbicides, the acetanilide herbicide may increase the retention and absorption of the other herbicide by the plant. Herbicide absorption can also be enhanced when in combination with oil-based adjuvants such as crop oil concentrate, free fatty acids, and esterified and saponified oils. Examples of such herbicides whose absorption is enhanced by oil-based adjuvants are cyclohexanidiones, aryloxyphenoxy, imidazolinone, and sulfonylurea herbicides.

The present invention is an improvement because it involves including in the herbicide spray solution, a material that modifies the surface properties of the spray solution, which results in decreased adherence of the spray droplets to the plant leaves. This is particularly useful when the plant is at a vulnerable stage of growth. For example, a crop such as dry beans can be at a sensitive growth stage (unifoliate) at the time it is most desirable to apply an acetanilide herbicide such as dimethamid. The present invention decreases foliar absorption of the herbicide

by the bean leaves, yet allows the herbicide to exert its action from the herbicide that has reached the soil.

Therefore, the present invention is a herbicide and water repellant composition wherein the surface properties of the composition is modified by the water repellant compound therein. Because these materials modify the surface properties of the herbicide spray solution, the spray droplets form spheres which bounce off the plant foliage. The water repellant compounds suitable for modifying the surface properties to make the herbicide composition of the present invention include an aqueous solution of an organosiliconate and aqueous siloxane solutions. An example of an aqueous solution of an organosiliconate that modifies the surface properties of the herbicide spray composition is an aqueous solution of sodium methyl siliconate. Sodium methyl siliconate has been used as a water repellant treatment for surfaces (see U.S. Patent No. 5,780,412 to Scarbrough and references therein). An example of an aqueous siloxane solution that is suitable is an aqueous solution of a water soluble silane coupling agent and an alkyltrialkoxysilane that modifies the surface properties of the herbicide spray composition such as an aqueous solution of N-(2-aminoethyl)-3-aminopropyltrimethoxysilane and methyltrimethoxysilane. When these compounds are mixed with an aqueous solution, they form an emulsion. Siloxanes are described in U.S. Patent 3,294,725 to Findlay et al. which is hereby incorporated herein by reference and aqueous solutions consisting of water soluble silane coupling agents and alkyltrialkoxysilanes are described in U.S. Patents 5,051,129 and 5,073,195, both to Cuthbert et al. which are hereby incorporated herein by reference to teach compositions which are suitable as repellant adjuvants in the present invention and methods for making them. While it may appear that any silicone containing

compound may be suitable for making the compositions of the present invention, the inventors have discovered otherwise. For example, methyltrimethoxysilane and a phosphonate ester alkyl silicon are water soluble silicon compounds. However, neither of these compounds is effective at producing a composition according to the present invention.

In an embodiment of the present invention disclosed herein, the herbicidal composition comprises isoxaflutole (5-cyclopropyl isoxazol-4-yl-2-mesyl-trifluoromethylphenyl ketone); a mixture of the herbicide metolachlor (2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide) or acetochlor, and the safener benoxacor ((4-dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine), MON 13900 or dichlormid; and any one of the repellant adjuvants disclosed herein. In a preferred embodiment, the herbicidal composition comprises the isoxaflutole as BALANCE, the metolachlor and benoxacor mixture as DUAL II, and the repellant adjuvant selected from the group consisting of an aqueous solution of sodium methyl siliconate and an aqueous solution of N-(2-aminoethyl)-3-aminopropyl-trimethoxysilane.

While the examples disclosed herein relate to the herbicide isoxaflutole, the present invention is not to be construed as being limited to the herbicide isoxaflutole. Examples of other herbicides which are encompassed by the present invention are nicosulfuron which is 2-[[[(4,6-dimethoxy-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]-N,N-dimethyl-3-pyridinecarboxamide - (ACCENT, Dupont, Wilmington, Delaware; isopropylamine salt, glyphosphate with adjuvants, (ACCORD, Monsanto Company, St. Louis, Missouri); primisulfuron which is methyl 2-[[[(4,6-bis(difluoromethoxy)-2-pyrimidinyl)amino]carbonyl]amino]sulfonyl]benzoate (BEACON, Novartis, Greensboro, North Carolina); Chlorimuron which is ethyl 2-[[[(4-chloro-

6-methoxy-2-pyrimidinyl)amino]carbonyl] amino]
sulfonyl]benzoate ethyl (CLASSIC, Du Pont, Wilmington,
Delaware); Glufosinate-ammonium salt which is (2-amino-
4-(hydroxymethylphosphinyl)butanoic acid (LIBERTY,
5 AgrEvo, Wilmington, Delaware); Linuron which is N1-(3,4-
dichlorophenyl)-N-methoxy-N-methylurea) (LOROX, Bayer,
Kansas City, Kansas); Linuron and chlorimuron ethyl
(LOROX PLUS, Dupont, Wilmington, Delaware);
Thifensulfuron which is (methyl 3-[[[(4-methoxy-6-
10 methyl-1,3,5-triazin-2-yl]carbonyl]amino] sulfonyl]-2-
thiophenecarboxylate) (PINNACLE, Dupont, Wilmington,
Delaware); Imazethapyr (PURSUIT, American Cyanamid,
Princeton, New Jersey); glyphosate-isopropyl amine salt
which is (N-(phosphonomethyl)glycine) (ROUNDUP, Monsanto
15 Company, St. Louis, Missouri); ROUNDUP with surface
components (phosphate esters and cationic tallow amines
(ROUNDUP ULTRA, Monsanto Company, St. Louis, Missouri);
imazaquin which is (2-[4,5-dihydro-4-methyl-4-(1-
methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinoline
20 carboxylic acid) (SCEPTER, American Cyanamid, Princeton,
New Jersey); acetochlor which is HARNESS and SURPASS
(available from Monsanto Company, and Zeneca Ag-Products
Wilmington, Delaware, respectively); alachlor, which is
2-chloro-2',6'-diethyl-N-(methoxymethyl)acetanilide and
25 sold as LASSO (available from Monsanto Company); EPTC
which is S-ethyl dipropylthiocarbonate and sold as
ERADICANE (available from Zeneca Ag-Products);
halosulfuron which is PERMIT and BATTALION (available
from Monsanto company); EPIC which is isoxaflutole, and
30 flufenacet (BAYFOE 5043) which is 4-fluoro-N-isopropyl-
2-[[5-trifluoromethyl)-1,3,4-thiadiazol-
2yl]oxy]acetamide (available from Bayer); and
glyphosphate-trimethylsulfonium salt (N-
(phosphonomethyl)glycine) (TOUCHDOWN, Zeneca Ag-
35 Products).

In addition to the composition of the present
invention containing a herbicide and the repellant

adjuvant, the present invention can further comprise activators, enhancers and safeners. Therefore, the present invention can further comprise a monosaccharide wherein the monosaccharide acts as an enhancer or potentiator for the herbicide in killing the weed without decreasing tolerance of the crop to the herbicide. Examples of such compositions are in U.S. Application Ser. No. 08/984,407 filed December 3, 1997 which is herein incorporated by reference. The present invention can further include oil-based adjuvants such as crop oil concentrate, free fatty acids, and esterified and saponified oils. The present invention can further include a safener which causes a reduction in injury to the crop plant without an unacceptable reduction in the herbicidal action. Examples of safeners encompassed by the present invention include benoxacor which is (4-dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine; dichlormid which is 2,2-dichloro-N,N-di-2-propenylacetamide (available from Zeneca, Inc.); MON 4660 which is available from Monsanto Company; R-29148 which is 2,2,5-trimethyl-N-dichloroacetyloxazolidine (available from Zeneca Ag-Products); R-25788 which is N,N-diallyl-2,2-dichloroacetamide (available from Zeneca Ag-Products); and MON 13900 which is 3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyl-oxazolidine (available from Monsanto Company). Other safeners include 2,2-dimethyl-5-phenyl-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-furanyl)-N-dichloroacetyl oxazolidine, 2,2-dimethyl-5(2-thienyl)-N-dichloroacetyl oxazolidine, 2,2-spirocyclohexy-N-dichloroacetyl oxazolidine, 4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1,4-benzoxazine, 3-[3-(dichloroacetyl)-2,2-dimethyl-5-oxalidinyl]pyridine, 4-(dichloroacetyl)-1-oxa-4-azapiro(4,5)-decane, 2,2-dichloro-1-(1,2,3,4-tetrahydro-1-methyl-2-isoquinolyl)ethanone, cis/trans-1,4-bis(dichloroacetyl)-2,5-dimethylpiperazine, N-

(dichloroacetyl)-1,2,3,4-tetrahydroquinaldine, 1,5-bis(dichloroacetyl)-1,5-diazacyclononane, 1-(dichloroacetyl)-1-azaspiro[4,4]nonane, and combinations thereof. Safeners are also disclosed in U.S. Patent 5,627,131 to Shribbs et al. which is hereby incorporated herein by reference. Examples of particular herbicide and safener combinations include DUAL II which consists of metolachlor and benoxacor (available from Novartis); SURPASS which consists of acetochlor and dichlormid (available from Zeneca Ag-Products); MON 8407 which consists of acetochlor and MON 4660 (available from Monsanto Company); ERADICANE which consists of EPTC and R-29148 (available from Zeneca Ag-Products); BATTALION which consists of a halosulfuron and MON 13900 (available from Monsanto Company); and MON 8411 which consists of acetochlor and MON 13900 (available from Monsanto Company). A particularly desirable combination of herbicide and safener is the herbicide acetochlor mixed with a safener selected from the group consisting of dichlormid, MON-13900 (flurilazole), R-29148, R-25788 (dichlormid), MON 4660 and combinations thereof.

The following examples are intended to promote a further understanding of the present invention.

EXAMPLE 1

Various silicone-based compounds were evaluated for ability to prevent herbicidal injury to the corn when used as repellent adjuvants in herbicide formulations. In previous experiments, it was shown that isoxaflutole in combination with metolachlor and benoxacor applied once corn had emerged caused significant corn injury. The basis of this severe injury is the retention and subsequent absorption of isoxaflutole in the foliar tissue of the corn plant, which is exacerbated by the metolachlor herbicide in the composition. Therefore, to address this problem, a number of silicon-based compounds, which were believed

to have repellant properties, were evaluated for the ability to prevent injury to corn plants. The compounds evaluated were an aqueous solution of 32 weight percent sodium methyl siliconate and 67 weight percent water ("SMS"), an aqueous solution of 35.7 weight percent N-(2-aminoethyl)-3-aminopropyltrimethoxysilane, 58.2 weight percent methyltrimethoxysilane, and 6.1 weight percent water ("AFS-MTMS"), methyltrimethoxysilane ("MTMS"), and phosphonate ester alkyl silicon ("PEAS"). These test adjuvants are commercially available from Dow Corning, Midland, Michigan. The test adjuvants were applied in combination with isoxaflutole or isoxaflutole in combination with metolachlor and the safener benoxacor, and the ability of the combination to reduce plant injury by inhibiting retention and subsequent absorption of the herbicide was evaluated.

Pioneer 3573 corn seeds (Pioneer 3573, a product of Pioneer Hi-Bred International, Inc., Des Moines, Iowa) were planted 2.54 cm deep, and velvetleaf and barnyardgrass seeds were planted 1.0 cm deep in 875 ml pots containing BACCTO professional greenhouse potting mix (a product of Michigan Peat Co., Houston, Texas). The seedlings were grown in a greenhouse maintained at 25°C \pm 2°C. Natural sunlight was supplemented with light from sodium vapor lamps, which provided a total midday light intensity of 1,000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photosynthetic photon flux at plant height during a 16 hour photoperiod. Plants were watered daily and fertilized weekly with 50 ml of a water soluble fertilizer solution (400 ppm nitrogen, 400 ppm P_2O_5 , and 400 ppm K_2O).

Isoxaflutole was combined with three different rates (0.25%, 0.5%, and 1.0%) of the four test adjuvants, which were then applied to postemergence to 2-leaf (5 inch) corn plants. Generally, corn leaf stages are described as the number of visible leaves. Isoxaflutole at 105 g/ha and isoxaflutole tank-mixed

with 1.1 kg/ha of metolachlor/benoxacor were applied either alone or in combination with ACTIVATOR 90, a non-ionic surfactant (NIS) product of Loveland Industries Inc., Greeley, Colorado) at 0.25% (v/v). The various herbicide applications were made to corn at the 2-leaf (V1) growth stage.

The herbicide compositions were applied through an 8003 E flat fan nozzle (available from Spraying Systems Co., Wheaton, Illinois) delivering 234 L/ha at a pressure of 172 kPa (25 gallons/acre). Corn tolerance was evaluated 8 days after treatment (DAT) by visually evaluating the plants for bleaching and necrotic symptoms and also by measuring corn height (base of the plant to its crown). Visual corn injury ratings were based on a scale from 0 to 100, with 0 indicating no effect and 100 indicating plant death. Corn height was measured in cm and presented as a percent of the non-treated plants, with 0 indicating total reduction in plant height and 100 indicating height equal to the non-treated plants. All experiments were conducted twice as completely randomized designs with four replications. Data were subjected to analysis of variance and means separated using Fisher's Protected LSD test at $\alpha=0.05$. Statistical analysis indicated no experimental run interactions, so the data were combined and reported as the means of two experiments. Non transformed means are presented since arcsine and square root transformations did not alter the interpretation of the data.

The data for the experiments are presented in Tables 1 and 2, which show that isoxaflutole when applied to corn by itself did not significantly injure the corn. But when isoxaflutole was applied in combination with metolachlor/benoxacor, severe corn injury of 47% occurred and plant height was reduced by 48%. However, when the mixture of isoxaflutole and metolachlor/benoxacor was mixed with either SMS or AFS-

MTMS, this injury was reduced to less than 10%. In contrast, neither of the other silicon-based compounds, MTMS or PEAS, reduced injury to corn when added to the isoxaflutole or isoxaflutole and metolachlor/benoxacor mixture. It is interesting that the other silicon-based compositions were not effective in ameliorating the herbicide's affect on the corn, in particular MTMS. Therefore, this example shows that only the adjuvants, SMS and AFS-MTMS, decreased the retention and subsequent absorption of isoxaflutole when either adjuvant was in combination with metolachlor/benoxacor.

5

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
22

Table 1.
Isoxaflutole Injury to Corn in Greenhouse Trials

Treatment	Adjuvant Rates			
	0%	0.25%	0.5%	1.0%
-----(% injury to corn)-----				
Isoxaflutole	2			
Isoxaflutole + Activator 90		15		
Isoxaflutole + PEAS		2	0	0
Isoxaflutole + SMS		0	0	0
Isoxaflutole + AFS-MTMS		0	0	0
Isoxaflutole + MTMS		1	0	0
Isoxaflutole + Metolachlor ^a	47			
Isoxaflutole + Metolachlor + Activator 90		59		
Isoxaflutole + Metolachlor + PEAS		52	49	51
Isoxaflutole + Metolachlor + SMS		8	9	7
Isoxaflutole + Metolachlor + AFS-MTMS		8	3	2
Isoxaflutole + Metolachlor + MTMS		51	56	56
^a The formulation metolachlor contained the herbicide safener benoxacor				

Table 2
Corn Height as a Percent of Control, 8 DAT in Green
House Trials

5	Treatment	Adjuvant Rates			
		0%	0.25%	0.5%	1.0%
		-----(% of control height)-----			
	Isoxaflutole	100			
	Isoxaflutole + Activator 90		96		
10	Isoxaflutole + PEAS		107	110	107
	Isoxaflutole + SMS		107	107	110
15	Isoxaflutole + AFS-MTMS		107	107	114
	Isoxaflutole + MTMS		103	110	107
	Isoxaflutole + Metolachlor ^a	62			
20	Isoxaflutole + Metolachlor + Activator 90		66		
25	Isoxaflutole + Metolachlor + PEAS		62	69	69
	Isoxaflutole + Metolachlor + SMS		100	100	100
30	Isoxaflutole + Metolachlor + AFS-MTMS		107	107	103
	Isoxaflutole + Metolachlor + MTMS		69	62	66
35	^a The formulation metolachlor contained the herbicide safener benoxacor				

The results of Tables 1 and 2 for the various
herbicide mixtures containing adjuvants added at a rate
of 0.5% are presented in Table 3. Table 3 shows that
injury to 2-leaf corn by isoxaflutole was reduced to 0
only when SMS or AFS-MTMS was mixed with the herbicide.
The table further shows that injury to corn caused by

isoxaflutole in combination with metolachlor/benoxacor was reduced from 47% to 8% and 3% by SMS and AFS-MTMS, respectively. The table also shows that isoxaflutole in combination with metolachlor/benoxacor reduced corn height by about 40% whereas adding either SMS or AFS-MTMS to the composition completely abrogated any affect the composition had on corn growth. The table clearly shows that the silicon-based compositions, PEAS and MTMS, were not useful as repellent adjuvants in herbicidal compositions. Therefore, this example demonstrates the present invention comprising either SMS or AFS-MTMS reduces herbicidal injury to corn when applied postemergence.

Table 3

Treatment ^a	Rate	Injury ^b		Height	
		Isoxaflutole ^c	Isoxaflutole + metolachlor ^d	Isoxaflutole	Isoxaflutole + metolachlor ^d
		-----%-----	-----%-----	-----% of control-----	-----% of control-----
Alone	0.25% v/v	2	47	100	64
+NIS	0.5% v/v	15	59	98	65
+PEAS	0.5% v/v	0	49	110	69
+SMS	0.5% v/v	0	8	106	99
+AFS-MTMS	0.5% v/v	0	3	107	105
+MTMS	0.5% v/v	0	56	108	62
LSD _{0.05}		-----4-----	-----8-----		

^aTreatments were applied to 2-leaf (12) corn

^bVisual injury ratings and corn heights were evaluated 8 DAT

^cIsoxaflutole was applied at 105 g/ha.

^dThe formulation of metolachlor contained the herbicide safener benoxacor and was applied at 1 l kg/ha

EXAMPLE 2

The greenhouse experiments showed that adding SMS or AFS-MTMS to a herbicide mixture rendered the herbicide safe for use on cultivated plants without reducing the herbicide's effective against weeds. However, greenhouse experiments are performed under controlled conditions. Therefore, field experiments were performed to assess how the present invention would

perform under actual farm conditions.

Conventional tillage experiments were conducted in 1998 to evaluate the influence of SMS and AFS-MTMS, on corn tolerance and weed control from postemergence applications of isoxaflutole alone and in tank-mixture with metolachlor/benoxacor. Experiments were conducted at the Michigan State University Crop and Soil Science Research Farm at East Lansing, Michigan on Capac sandy clay loam soil (fine-loamy, mixed mesic Acric Ochraqualfs) containing 3.1% organic matter with a pH of 6.3 in 1998.

Tillage consisted of moldboard plowing in the fall prior to spring disking and field cultivation. Prior to spring cultivation, 320 kg/ha of 46-0-0 fertilizer was applied broadcast. At planting, 140 kg/ha of 6-24-24 fertilizer was applied as a banded treatment 5 cm below and 5 cm beside the corn seeds. Pioneer 37R71 corn was planted on May 11, 1998 at a rate of 62,000 seeds/ha. Each plot was 10.6 m long and consisted of 4 rows spaced 76 cm apart.

The tillage experiments were conducted as a randomized complete block design in a factorial arrangement with three replications. The factors consisted of herbicide application timing and herbicide treatment. Herbicides were applied when the corn was at the 2-leaf and 4-leaf stages. Corn leaf stages are described as the number of visible leaves. Herbicide treatments included isoxaflutole alone (BALANCE) at 105 g/ha and in combination with 1.1 kg/ha of metolachlor/benoxacor (DUAL II). Each of these treatments were applied either alone or with either SMS or AFS-MTMS. Each repellent was used at a rate of 0.5% v/v. Additional treatments not included in the factorial arrangement were an untreated check and a weed-free check. All herbicides were applied with a tractor mounted, compressed-air sprayer calibrated to deliver 2.6 L/ha at 207 kPa using 8003 E flat-fan

nozzles.

Corn tolerance was evaluated 30 days after planting (DAP) by visually evaluating plants for bleaching and necrotic symptoms and also by measuring corn height (base of plant to the crown) 40 DAP. Weed control by species was visually evaluated 60 DAP. Visual evaluations were based on a scale of 0 (no effect) to 100% (complete weed or crop death). Corn grain yield was determined by harvesting the center two rows of each plot with a plot combine. Seed weight was adjusted to 15% moisture.

Data were subjected to analysis of variance and means separated using Fisher's Protected LSD test at $\alpha=0.05$. Data were combined over years when treatment and/or application timing by year interactions were not significant $\alpha=0.05$. Non-transformed means for corn injury and weed control are presented since arcsine and square root transformations did not alter the interpretation of the data. Corn height and yield results were converted to a percent of the weed-free treatment after separation.

Herbicide application times, corn stages, weed heights, and densities for the field trial in 1998 are presented in Table 4 and rainfall data in Table 5.

Table 4

	1998	
	2-Leaf ^a	4-Leaf
Days after planting ^b	9	15
Corn		
Leaves with collars	1	2
Ave height (cm)	10	13
Giant foxtail		
Ave. height (cm)	0.6	4
Density (plants/m ²)	33	65
Broadleaf weeds ^c		
Ave. height	0.6	3
Ave density (plants/m ²)	44	44

^a Corn leaf stage refers to the number of visible leaves.

^b Corn planted May 11, 1998

^c Broadleaf weeds include: common lambsquarters, redroot pigweed, common ragweed, and velvetleaf.

Table 5

Days after planting	Amount of Rainfall
	1998
	-----mm-----
0-7	2
8-14	3
15-21	12
22-28	0
Total	17

Corn (ZEAMX) in the 2-leaf or 4-leaf stage was treated with herbicidal compositions consisting of BALANCE (isoxaflutole); BALANCE and DUAL II (metolachlor and the safener benoxacor); BALANCE and SMS; BALANCE and AFS-MTMS; BALANCE, DUAL II and SMS; or BALANCE, DUAL, and AFS-MTMS. The percent injury was determined 6 DAT, 12 DAT, and 30 DAP. As weed controls, the annual grasses (ANGR) and common lambsquarters, *Chenopodium album* L. (CHEAL) were treated with herbicidal compositions consisting of BALANCE; BALANCE and DUAL II; BALANCE and SMS; BALANCE and AFS-MTMS; BALANCE, DUAL II and SMS; or BALANCE, DUAL, and AFS-MTMS. The percent injury for the weed controls was determined 30 DAP.

The results are shown in Table 6. The results show that the repellent adjuvants SMS and AFS-MTMS were effective in reducing the percent injury to corn caused by the herbicide when either was included in herbicidal compositions consisting of BALANCE and DUAL II. AFS-MTMS was particularly effective, when it was included in the herbicidal composition and applied to 2-leaf corn plants, corn injury 6 DAT was only 7.3%, whereas without AFS-MTMS, the corn injury was 65%. By 12 DAT and beyond, corn injury of caused by the BALANCE and DUAL II composition containing AFS-MTMS was not detectable whereas without AFS-MTMS, the injury remained about 55-50%. The weed controls show that SMS and AFS-MTMS do not appear to reduce the efficacy of BALANCE and DUAL II

to control weeds.

The results in Table 6 also show that SMS and AFS-MTMS reduced injury to corn at the 4-leaf stage when included in herbicidal compositions containing both BALANCE and DUAL II. In particular, the percent injury to corn was significantly reduced when the herbicidal composition included AFS-MTMS as the repellent adjuvant. The results further show that including a safener in the herbicide composition (the benoxacor) had no safening effect when used in combination with isoxaflutole. These results demonstrate that the present invention is useful and effective under actual field conditions.

5.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200

Table 6

Weed Code: Crop Code:						ZEAMX			ANGR	CHEAL
Treatment/Evaluation Interval Date evaluated						6 DAT 5-26-98	12 DAT 6-1-98	30 DAP 6-9-98	30 DAP 6-9-98	30 DAP 6-9-98
No.	Composition	Form Amt	Rate	Rate Unit	Grow Stg	Injury Percent			Control Percent	
1	BALANCE	75	1.5	OZ A/A	2-leaf	0.0	0.0	0.0	41.7	98.3
2	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	0.0	0.0	0.0	50.0	100.0
3	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	0.0	0.0	0.0	35.0	97.3
4	BALANCE DUAL II	75 7.8	1.5	OZ A/A LB A/A	2-leaf	65.0	55.0	50.0	99.3	100.0
5	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	23.3	5.0	5.7	96.0	100.0
6	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	7.3	0.0	0.0	98.7	100.0
7	BALANCE	75	1.5	OZ A/A	4-leaf		16.7	9.7	48.3	100.0
8	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	4-leaf		15.0	10.7	41.7	100.0
9	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	4-leaf		10.0	10.0	55.0	48.3
10	BALANCE DUAL II	75 7.8	1.5 1.0	OZ A/A LB A/A	4-leaf		81.7	83.3	100.0	100.0
11	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf		48.3	30.0	100.0	99.3
12	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf		26.7	16.7	98.7	99.7
13	Untreated						0.0	0.0	0.0	0.0
LSD (P= 0.05)						4.65	7.99	7.52	14.50	15.82
Standard Deviation						2.61	4.74	4.46	8.61	9.39
CV						19.11	23.87	26.86	12.94	10.68

BALANCE - isoxaflutole; DUAL II - metolachlor/benoxacor ; DAT - day after treatment, DAP - day after planting; ZEAMX - corn, *Zea mays* L.; ANGR - annual grasses, CHEAL - common lambsquarter, *Chenopodium album* L.

Additional weed controls consisted of the weeds giant foxtail, *Setaria faberi* Herrm. (SETFA);

10

Table 7

						Weed Code:		
						AMARE	AMBEL	ABUTH
Treatment/Evaluation Interval						30 DAP	30 DAP	30 DAP
Date evaluated						6-9-98	6-9-98	6-9-98
No.	Composition	Form Amt	Rate	Rate Unit	Grow Stg	Control Percent		
1	BALANCE	75	1.5	OZ A/A	2-leaf	100.0	100.0	98.3
2	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	100.0	100.0	99.3
3	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	2-leaf	95.0	100.0	100.0
4	BALANCE DUAL II	75 7.8	1.5	OZ A/A LB A/A	2-leaf	100.0	100.0	100.0
5	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	100.0	100.0	97.7
6	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	2-leaf	100.0	100.0	99.3
7	BALANCE	75	1.5	OZ A/A	4-leaf	100.0	100.0	95.7
8	BALANCE SMS	75	1.5 0.5	OZ A/A % V/V	4-leaf	100.0	100.0	97.0
9	BALANCE AFS-MTMS	75	1.5 0.5	OZ A/A % V/V	4-leaf	98.3	97.3	95.7
10	BALANCE DUAL II	75 7.8	1.5 1.0	OZ A/A LB A/A	4-leaf	100.0	100.0	98.3
11	BALANCE DUAL II SMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf	100.0	100.0	97.0
12	BALANCE DUAL II AFS-MTMS	75 7.8	1.5 1.0 0.5	OZ A/A LB A/A % V/V	4-leaf	100.0	100.0	98.3
13	Untreated					0.0	0.0	0.0
LSD (P= 0.05)						4.16	2.16	2.89
Standard Deviation						2.47	1.28	1.72
CV						2.69	1.39	1.9

BALANCE - isoxaflutole, DUAL II - metolachlor/benoxacor, DAT - day after treatment; DAP - day after planting, AMARE -redroot pigweed, *Amaranthus retroflexus* L., AMBEL - common ragweed, *Ambrosia artemisiifolia* L., ABUTH - velvetleaf, *Abutilon theophrasti* Medicus

The results of Tables 6 and 7 are condensed in Table 8, which shows that the present invention prevented injury to corn caused by herbicide component

of the composition without substantially affecting the ability of the herbicide component to control weed growth. Additional data was added to Table 8. The additional data shows that adding SMS or AFS-MTMS to the herbicide mixture prevented the herbicide component from affecting the growth of the corn. Also new in Table 8 is data showing that the herbicide mixture containing either SMS or AFS-MTMS did not substantially alter the herbicide component's ability to control the growth of giant foxtail (SETFA). Significantly, the grain yield of corn treated with the present invention was greater than the yields of corn treated with the herbicide mixture lacking either SMS or AFS-MTMS or the untreated controls.

Therefore, the results of the field trial show that the present invention renders the herbicides comprising the invention safe for use on corn while not reducing the ability of the herbicides comprising the present invention to control a wide variety of weeds.

Table 8

Herbicide	Application Stage	Corn		Weed Control ^c				Grain Yield
		Injury ^a	Height ^b	SETFA	CHEAL	AMARE	ABUTH	
		—%—	—cm—	—%—				kg/ha
Isoxaflutole ^d	2-leaf	0	61	73	95	98	99	10420
Isoxaflutole + SMS ^e	2-leaf	0	61	72	96	98	100	10146
Isoxaflutole + AFS-MTMS	2-leaf	0	61	56	85	92	100	10146
Isoxaflutole + Metolachlor ^g	2-leaf	50	41	95	100	100	100	9617
Isoxaflutole + metolachlor + SMS	2-leaf	6	56	91	100	97	98	10277
Isoxaflutole + metolachlor + AFS-MTMS	2-leaf	0	62	86	98	100	97	10527
Isoxaflutole	4-leaf	10	61	70	96	100	100	9995
Isoxaflutole + SMS	4-leaf	11	61	63	100	100	100	9881
Isoxaflutole + AFS-MTMS	4-leaf	10	58	68	83	95	96	10728
Isoxaflutole + Metolachlor ^a	4-leaf	83	32	95	100	100	98	9192
Isoxaflutole + metolachlor + SMS	4-leaf	30	50	99	100	100	100	9508
Isoxaflutole + metolachlor + AFS-MTMS	4-leaf	17	58	96	100	100	100	9993
Untreated		0	64	0	0	0	0	8528
LSD ₅₀		8	7	13	9	8	3	850

^a Corn injury evaluated 30 DAP which was 21 days after 2-leaf application and 15 days after 4-leaf application

^b Corn height was measured 40 DAP.

^c Weed control was evaluated 60 DAP.

^d The rate of isoxaflutole was 105 g/ha

^e All adjuvants were applied at 0.5% v/v.

^f The rate of metolachlor was 1121 g/ha

^g The formulation of metolachlor contained the herbicide safener benoxacor

EXAMPLE 3

This example was performed as in Example 1, except that adjuvants SMS and AFS-MTMS were added to an isoxaflutole tank-mixture that contained an acetochlor herbicide and the safener MON-13900. As in Example 1, the herbicide applications were made to 2-leaf corn, and the herbicide treatments consisted of isoxaflutole at 105 g/ha and isoxaflutole tank-mixed with 1.8 kg/ha of acetochlor/MON-13900 applied either alone or in combination with either NIS at 0.25% (v/v), SMS at 0.5% (v/v), or AFS-MTMS at 0.5% (v/v).

The herbicide compositions were applied through an 8003 E flat fan nozzle delivering 234 L/ha at a pressure of 172 kPa. Corn tolerance was evaluated 8 DAT by visually evaluating the plants for bleaching and necrotic symptoms and also by measuring corn height (base of the plant to its crown). Visual corn injury ratings were based on a scale from 0 to 100, with 0 indicating no effect and 100 indicating plant death. Corn height was measured in cm and presented as a percent of the non-treated plants, with 0 indicating total reduction in plant height and 100 indicating height equal to the non-treated plants. All experiments were conducted twice as completely randomized designs with four replications. Data were subjected to analysis of variance and means separated using Fisher's Protected LSD test at $\alpha=0.05$. Statistical analysis indicated no experimental run interactions, so the data were combined and reported as the means of two experiments. Non transformed means are presented since arcsine and square root transformations did not alter the interpretation of the data.

Table 9 shows the corn injury and height reductions as influenced by SMS or AFS-MTMS combined with isoxaflutole alone and in combination with acetochlor/MON-13900 when applied postemergence in the greenhouse. The table shows that the percent injury to

corn treated with isoxaflutole in combination with acetochlor/MON-13900 was reduced to 20% when SMS was added to the combination. When AFS-MTMS was added to the combination of isoxaflutole and acetochlor/MON-13900 there was no visible injury to the corn. The table also shows that isoxaflutole in combination with acetochlor/MON-13900 reduced corn height by about 50% whereas adding either SMS or AFS-MTMS to the composition reduced the effect of the composition on corn height. In particular, in the presence of SMS, the height of the corn was reduced by only 20% and in the presence of AFS-MTMS, the height of the corn was virtually unaffected.

Table 9

Treatment ^a	Rate	Injury ^b		Height	
		Isoxaflutole ^c	Isoxaflutole ^c + acetochlor ^d	Isoxaflutole	Isoxaflutole ^c + acetochlor ^d
		-----%		-----% of control-----	
Alone		0	64	99	47
+NIS	0.25% v/v	13	68	88	45
+SMS	0.5% v/v	0	20	100	79
+AFS-MTMS	0.5% v/v	0	0	100	98
LSD _{0.05}		-----4-----		-----8-----	

^a Treatments were applied to 2-leaf (12) corn.

^b Visual injury ratings and corn heights were evaluated 8 DAT.

^c Isoxaflutole was applied at 105 g/ha.

^d The formulation of acetochlor contained the herbicide safener MON-13900 and was applied at 1.1 kg/ha

EXAMPLE 4

This example was conducted to determine the effect of SMS and AFS-MTMS on foliar retention of isoxaflutole either alone or in combination with metolachlor/benoxacor. A version of the technique reported by Boldt and Putnam, Weed Science 28: 474-477 (1980) was used. Herbicide treatments examined were isoxaflutole applied alone at 105 g/ha and isoxaflutole tank-mixed with 1.1 kg/ha of metolachlor/benoxacor.

Each of these herbicide treatments were applied alone or with either SMS or AFS-MTMS. The repellent adjuvants were each used at a rate of 0.5% v/v. The spray treatments, which included CHICAGO SKY BLUE (a product available from Sigma Chemical Co., St. Louis, Missouri) at 2.5 g/L, were applied to 2-leaf corn. Immediately after application, the whole plant was harvested and rinsed with distilled water containing the non-ionic surfactant X-77 (a product available from Valent U.S.A. Corp., Walnut Creek, California) at 0.25% v/v. The absorbance of the rinsate was determined spectrophotometrically at 625 nm. Dye retention ($\mu\text{g}/\text{plant}$) was calculated from a standard curve.

Table 10 shows the spray retention of isoxaflutole as influenced by either SMS or AFS-MTMS, alone and in combination with metolachlor/benoxacor applied to 2-leaf corn in the greenhouse. The results show that neither SMS or AFS-MTMS increased the retention of isoxaflutole alone by the corn plant. The results further show that AFS-MTMS was particularly effective in reducing isoxaflutole retention when the composition further included metolachlor/benoxacor.

Table 10

Treatment	Rate	Spray retention	
		Isoxaflutole ^a	Isoxaflutole + metolachlor ^b
-----μg of isoxaflutole/plant-----			
Alone		4.4	15.1
+SMS	0.5% v/v	5.8	17.3
+AFS-MTMS	0.5% v/v	4.4	6.2
LSD _{0.05}		-----2.2-----	

^a rates; isoxaflutole at 105 g/ha, metolachlor/benoxacor at 1.1 kg/ha.

^b The formulation contained the herbicide safener benoxacor

EXAMPLE 5

This example was to determine whether the adjuvants NIS, MTMS, SMS, AFS-MTMS, and PEAS affected weed control from early postemergence applications of isoxaflutole and isoxaflutole tank-mixed with metolachlor/benoxacor. Reduced rates of isoxaflutole (53 g/ha) and isoxaflutole tank-mixed with metolachlor/benoxacor (0.55 kg/ha) were applied alone and with either of NIS, MTMS, SMS, AFS-MTMS, or PEAS to 2-leaf velvetleaf (ABUTH) (3.5 to 5 cm) and 3-leaf barnyardgrass (5 to 10 cm). Velvetleaf and barnyardgrass controls were evaluated 21 DAT and shoots of both species were harvested to measure dry weight per pot. The results of this example are consistent with the weed control results presented in Tables 6, 7 and 8 of Example 2, which showed that the present invention was effective against weeds.

While the present invention is described herein with reference to illustrated embodiments, it should be understood that the invention is not limited hereto. Those having ordinary skill in the art and access to the teachings herein will recognize additional modifications and embodiments within the scope thereof. Therefore, the present invention is limited only by the Claims attached herein.